## SOLAR OBSERVATIONS.

## SOLAR AND SKY RADIATION MEASUREMENTS DURING AUGUST, 1923.

By IRVING F. HAND, Acting in Charge, Solar Radiation Investigations.

For a description of instruments and exposures, and an account of the method of obtaining and reducing the measurements, the reader is referred to the Review for April, 1920, 48:225, and a note in the Review for November, 1922, 50:595.

From Table 1 it is seen that direct solar radiation intensities averaged below the normal values for August at Washington, D. C., and above normal at Madison, Wis., and Lincoln, Nebr. A noon intensity of 1.46 gram-calories per minute per square centimeter measured on the 24th at Madison, Wis., is the highest ever measured at that station in August.

Table 2 shows deficiencies in the radiation received from the sun and sky at all three stations, the deficiency being especially marked during the first week at

Washington.

Skylight polarization measurements obtained at Washington on 11 days give a mean of 47 per cent with a maximum of 58 per cent on the 14th, At Madison measurements on 7 days give a mean of 66 per cent with a maximum of 74 per cent on the 19th. These are below average values for August at Washington, but decidedly above normal at Madison. In fact, the value obtained on the 19th is the highest ever obtained at Madison in August.

Table 1.—Solar radiation intensities during August, 1923.

[Gram-calories per minute per square centimeter of normal surface.]

Washington, D. C.

	Sun's zenith distance.											
	8 a.m.	78.7°	75.7°	70.7°	60.0°	0.0°	60.0°	70.7°	75.7°	78.7°	Noon	
Date.	75th mer. time.	Air mass.									Local	
		А. М.					Р. М.					
		5.0	4.0	3.0	2.0	*1.0	2.0	3.0	4.0	5.0	е.	
Aug. 13	mm. 15.65	cal. 0.55	cal. 0.68	cal. 0.83	cal. 1.07	cal. 1.19	cal. 0.71 0.98	cal.	cal.	cal.	mm. 11.31	
14 15 16 20	. 10. 21 . 17. 87 . 11. 81 . 13. 61	0.61	0.71	0.83	0.99 0.63	1.27 1.19 0.90 1.09	0.98				11.3 12.6 11.8	
21 22 24	16. 20 9. 47 10. 97					1.20	1.20				17.9 8.1 7.8	
25 30 31	. 12. 68 . 13. 61 . 15. 11		0.40	0. 55	0.98 0.81 0.55	1.32 1.00 0.82					9.4 16.2 12.6	
Means Departures		(0. 58) 0. 04										

<sup>\*</sup> Extrapolated.

Table 1.—Solar radiation intensities during August, 1923—Continued.

Madison, Wis.

	Sun's zenith distance.										
	8 a.m.	18.7°	75.7°	70.7°	60.0°	0.0°	60,0°	70.0°	75.7°	18.7°	Noon
Date.	75th mer. time.	Air mass.									Loca mea
		А. М.					Р. М.				solar time
		5.0	4.0	3.0	2,0	١ 1.0	2,0	3,0	4.0	5,0	e.
Aug. 17	mm. 10.59	cal.	cal.	cal.	cal. 1.00	cal. 1.28	cal.	cal.	cal.	cal.	mm. 10. 9
18.: 19 22	10. 97 10. 97 7. 04				1.22 1.20	1.80 1.40					11.8 10.2 7.2
23 24 29	7.87 7.29 9.14			0.90 1.09	1.13 1.31 1.06	1.41 1.49	1.13 1.30				6. 5 10. 5 10. 5
30 Means	12.68		 	(1.00)			(1.22)				14.6
Departures				+0.07	+0.00	+0.02	+0.18			<b></b> -	

Lincoln, Nebr.											
Aug. 10 11	17.37 15.65 14.10		0.80	0.97	1. 16	1.35	1.01	0.91 1.01	0. 79 0. 88	0.69	18. 59 17. 37 13. 13
13 14 22	14.88 17.37 7.04		0.83 0.93	1.02 0.92 1.05	1.21 1.13 1.22	1.45 1.27	1.13	0.93	0.79	0.69	14. 10 20. 57 7. 29
24 25	9. 14 1. 38		0.74	1.01			1.02				8. 48 13. 13
Means Departures			0.82 +0.03	0. 99 +0. 09		1.36 +0.08				(0. 69) —0. 02	

TABLE 2.—Solar and sky radiation received on a horizontal surface.

Week	Average	daily ra	diation.	Average for	daily de	eparture k.	Excess or deficiency since first of year.			
beginning.	Wash- ington.	Madi- son.	Lin- coln.	Wash- ington.	Madi- son.	Lin- coln.	Wash- ington.	Madi- son.	Lin- coln.	
July 30 Aug. 6 13 20 27	cal. 279 395 451 487 406	cal. 438 420 492 427 291	cal. 462 388 489 530 439	cal. -188 -61 +13 +69 -1	cal. -34 -36 +51 +4 -111	cal. -73 -128 -4 +55 -19	cal. -3,360 -3,788 -3,694 -3,213 -3,223	cal. +676 +424 +780 +811 +34	cal. -1,23 -2,12 -1,15 -1,77 -1,90	